

602600

FTD-MT- 63-194

# TRANSLATION

MECHANISM OF DIRECT ACTION OF RADIATION ON PERCHLORIC U.C. ACID

By



L. T. Bugayenko

## FOREIGN TECHNOLOGY DIVISION

AIR FORCE SYSTEMS COMMAND

WRIGHT-PATTERSON AIR FORCE BASE

OHIO



## FOREWORD

This document is a machine translation of Russian text which has been processed by the AN/GSQ-16(XW-2) Machine Translator, owned and operated by the United States Air Force. The machine output has been fully post-edited. Ambiguity of meaning, words missing from the machine's dictionary, and words out of the context of meaning have been corrected. The sentence word order has been rearranged for readability due to the fact that Russian sentence structure does not follow the English subject-verb-predicate sentence structure. The fact of translation does not guarantee editorial accuracy, nor does it indicate USAF approval or disapproval of the material translated.

193854

**CLEARINGHOUSE FOR FEDERAL SCIENTIFIC AND TECHNICAL INFORMATION CFSTI  
DOCUMENT MANAGEMENT BRANCH 410.11**

**LIMITATIONS IN REPRODUCTION QUALITY**

**ACCESSION #**

*AD 602600*

- ☒ 1. WE REGRET THAT LEGIBILITY OF THIS DOCUMENT IS IN PART UNSATISFACTORY. REPRODUCTION HAS BEEN MADE FROM BEST AVAILABLE COPY.
- ☐ 2. A PORTION OF THE ORIGINAL DOCUMENT CONTAINS FINE DETAIL WHICH MAY MAKE READING OF PHOTOCOPY DIFFICULT.
- ☐ 3. THE ORIGINAL DOCUMENT CONTAINS COLOR, BUT DISTRIBUTION COPIES ARE AVAILABLE IN BLACK-AND-WHITE REPRODUCTION ONLY.
- ☐ 4. THE INITIAL DISTRIBUTION COPIES CONTAIN COLOR WHICH WILL BE SHOWN IN BLACK-AND-WHITE WHEN IT IS NECESSARY TO REPRINT.
- ☐ 5. LIMITED SUPPLY ON HAND: WHEN EXHAUSTED, DOCUMENT WILL BE AVAILABLE IN MICROFICHE ONLY.
- ☐ 6. LIMITED SUPPLY ON HAND: WHEN EXHAUSTED DOCUMENT WILL NOT BE AVAILABLE.
- ☐ 7. DOCUMENT IS AVAILABLE IN MICROFICHE ONLY.
- ☐ 8. DOCUMENT AVAILABLE ON LOAN FROM CFSTI ( TT DOCUMENTS ONLY).
- ☐ 9.

**NBS 9/64**

**PROCESSOR:**

*BL*

COMPLETE FOR NASA CONTRACTOR DOCUMENTS ONLY

DOCUMENT TYPES

1. \_\_\_\_\_ Report is unclassified and is suitable for unlimited announcement and distribution to the aeronautics and space community, and for public sale.
2. \_\_\_\_\_ Report is unclassified, but contains information of limited usefulness and does not justify widespread automatic distribution to the aeronautical and space community. It can, however, be announced and made publicly available.
3. \_\_\_\_\_ Report is unclassified, but for official reasons, must be restricted as follows:
  - a. \_\_\_\_\_ Government agencies and their contractors only.
  - b. \_\_\_\_\_ Government agencies only.
  - c. \_\_\_\_\_ NASA activities only.
4. \_\_\_\_\_ Report bears a security classification and is suitable for distribution within the limits of security considerations to:
  - a. \_\_\_\_\_ Government agencies and their contractors only.
  - b. \_\_\_\_\_ Government agencies only.
  - c. \_\_\_\_\_ NASA activities only.
5. \_\_\_\_\_ Reprint of a journal article reporting NASA-supported work.
6. \_\_\_\_\_ Article prepared for journal publication (preprint or other copy) reporting NASA-supported work. (Normally handled as No. 2 above.)

Estimated date of publication: \_\_\_\_\_

7. \_\_\_\_\_ Material for presentation at a meeting or conference  
Name of Meeting: \_\_\_\_\_ Date: \_\_\_\_\_  
Sponsor(s): \_\_\_\_\_

- a. \_\_\_\_\_ Scheduled for publication in proceedings. (Normally handled as No. 2 above.)

Estimated date of publication: \_\_\_\_\_

Not scheduled for publication in proceedings and subject to the following limitations or announcement and dissemination:

- b. \_\_\_\_\_ Government agencies and their contractors only.
- c. \_\_\_\_\_ Government agencies only.
- d. \_\_\_\_\_ NASA activities only.

602600

# SCIENTIFIC AND TECHNICAL INFORMATION FACILITY

operated for National Aeronautics and Space Administration by Documentation Incorporated

## LOAN DOCUMENT TRANSMITTAL

Please attach a copy of this form to any loan documents sent to the Scientific and Technical Information Facility. A single copy of this form and the document should be sent to the following address:

Scientific and Technical Information Facility  
Attention: NASA Representative  
Box 5700  
Bethesda 14, Maryland

Every effort will be made to return this document in the shortest possible time.

Please complete.

Sent by:

**SCIENTIFIC AND TECHNICAL**  
INFORMATION FACILITY  
POST OFFICE BOX 5700  
BETHESDA, MD. 20014

Date sent: \_\_\_\_\_

☐

checked for N numbers

☐

checked against N-62 numbers

See other side for NASA contractor report

.....  
For Facility Use Only

## LOAN DOCUMENT PROCESSING RECORD

N62- \_\_\_\_\_

Date received: \_\_\_\_\_

X62- \_\_\_\_\_

### THIS DOCUMENT IS ON LOAN

Please process as follows:

☐

Film for Microform and make Xerox copy

Date

Initials

☐

Microform completed

☐

Xerox copy made

☐

Xerox copy sent to Input Station

☐

Original sent to Mail Room

☐

Document returned to sender

7428

THIS LOAN DOCUMENT FORM MUST BE SENT TO THE CASE FILE

# EDITED MACHINE TRANSLATION

MECHANISM OF DIRECT ACTION OF RADIATION ON PERCHLORIC U.C.  
ACID

BY: L. T. Bugayenko

English Pages: 9

SOURCE: Trudy II Vsesoyuznogo Soveshchaniya Po Radiatsionnoy  
Khimii, Izdatel'stvo AN SSSR, Moskva, 1962, pp 144-148

S/2844/62/0/0

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:




TRANSLATION DIVISION  
FOREIGN TECHNOLOGY DIVISION  
WP-APB, OHIO.

Tape No. MT-63-194  
 Transactions of the Second All-Union  
 Conference on Radiation Chemistry,  
 Moscow, 1962.  
 Pages 144-148


# MECHANISM OF DIRECT ACTION OF RADIATION ON PERCHLORIC U.C. ACID

L. T. Bugayenko

Earlier it was shown that the perchlorate ion is reduced only as a result of direct action of radiation [1,2]. Neither atoms of hydrogen nor hydroxyl radicals interact with perchlorate ion. The mechanism of direct action appeared complicated, proceeding through several intermediate steps [3]:



Excited perchlorate, ions most probably, dissipate their energy during collisions, but to a certain extent the disintegration of excited ions by the reaction




is possible.

In this work we continued investigation of the mechanism of transformation of perchlorate, ions studying the influence of ions of bivalent iron and molecules of organic compounds (ethyl alcohol and acetone) on yields of reduction of perchlorate ion and yields of reduction products of perchlorate ions.


Preparation, experiment, and analyses were conducted as described in [2,3]. Irradiation was produced by x-rays with maximum energy of 70 kv.

It was found that in oxygen saturated solutions of iron perchlorate in 2.5 M  $\text{HClO}_4$ , the yield of reduction of perchlorate ion in the interval of concentrations of  $\text{Fe}^{2+}$  ion, 0.005 - 0.06 M, maintains a constant value of  $1.20 \pm 0.05$  ions per 100 ev of absorbed energy. It was found that the yields of formation of ions of chlorate and chloride depend on the concentration of  $\text{Fe}^{2+}$  ion. These data are shown in Fig. 1. The observed dependence of  $G(\text{ClO}_3^-)$  on  $[\text{Fe}^{2+}]$  shows that in the formation of chlorate ion reaction (3), not reaction (5) plays the main role, since the yield of formation of chlorate as a result of disintegration of excited perchlorate ions should not depend on the concentration of dissolved substances.

In Fig. 2 is shown a change of the ratio  $G(\text{ClO}_3^-)/G(\text{Cl}^-)$  with increase of concentration of iron. Such a sharp drop of the curve is connected, first of all, with competition between dioxide of chlorine and the  $\text{Fe}^{2+}$  ion for radical  $\text{OH}$  and, secondly, with the very fast reaction:



The formed ion of chlorite quickly is reduced by iron ions to the chlorine ion:



In Fig. 3 is shown the change of the yield of formation of  $\text{Fe}^{3+}$  ion as a function of concentration of  $\text{Fe}^{2+}$  ion.




Fig. 1. Dependences of the yield of disintegration of ion of perchlorate (1), of yields of formation of ion of chlorate (2) and of ion of chloride (3) on the concentration of perchlorate of bivalent iron in 2.5 M  $\text{HClO}_4$ . KEY: (a) G, ions/100 ev; (b) Concentration of  $\text{Fe}^{2+}$  ions,  $\text{M} \cdot 10^3$ .







Fig. 2. Dependence of ratio of yields of ions of chlorate and ions of chlorine on the concentration of iron in 2.5 M HClO<sub>4</sub>. KEY: (a) Concentration of Fe<sup>2+</sup> ions, M · 10<sup>2</sup>.

Experimental points lie on a straight line. If we extrapolate this straight line to the y-axis, we will obtain the value of yield of oxidation of iron under conditions when it has practically no influence on the yield of reduction products of ion of perchlorate. The extrapolated value of yield of oxidation of iron equals  $17 \pm 1$  ion per 100 ev. We will make an attempt to calculate the value of yield of oxidation of ion of iron in these conditions, assuming the known values of yield of products of radiolysis of water. We will consider that the yield of products of transformation of water are the same as in 0.4 M H<sub>2</sub>SO<sub>4</sub> [4]. The electronic fraction of water in 2.5 M HClO<sub>4</sub> equals 0.8; therefore we will have the following values of yield  $G_H = 3.00$ ;  $G_{OH} = 2.50$  and  $G_{H_2O_2} = 0.60$ . The electron appearing during direct action of radiation on the ion of perchlorate, according to Lee [5], will cause formation of one H radical:



As was shown above, the fraction of the reaction (5) is small; therefore for calculation we will assume that 1.25 electrons will be formed and, consequently, 1.25 additional atoms of hydrogen per 100 ev. Radical OH oxidizes ions of iron and dioxide of chlorine. For the fraction of ions of iron will be obtained  $G_{OH} - G(ClO_3^-) = 1.50$  radicals of OH per 100 ev. Since there is oxygen in solution, the atoms of hydrogen will be converted into radicals of HO<sub>2</sub>. The latter very effectively interact with ions of Fe<sup>2+</sup>, the concentration of which is significantly greater than the concentration of dioxide of chlorine.

Therefore in the presence of ions of iron the reduction of  $\text{ClO}_2$  will be produced not by radicals  $\text{HO}_2$ , but by ions of  $\text{Fe}^{2+}$  by reactions (6) and (7).




Fig. 3. Dependence of yield of oxidation of iron on the concentration of perchlorate of iron in 2.5 M  $\text{HClO}_4$ .  
KEY: (a)  $G(\text{Fe}^{3+})$ , ions/100 ev. (b) Concentration of ions  $\text{Fe}^{2+}$ , M  $\text{HClO}_4$ .

The yield of formation of ions of chlorine equals 0.22 ions per 100 ev. Full yield of oxidation of ions of iron will be equal to:

$$G(\text{Fe}^{3+}) = 3G_{\text{H}} + 2G_{\text{H}_2\text{O}_2} + G_{\text{OH}} - G(\text{ClO}_2) + 5G(\text{Cl}^-). \quad (9)$$

Substituting the above calculated values of yields, we obtain  $G(\text{Fe}^{3+}) = 16.5$  ions per 100 ev. This magnitude is in good agreement with the value of yield of oxidation of iron found by extrapolation of experimental data




Fig. 4. Dependence of yield of disintegration of ion of perchlorate on the concentration of ethyl alcohol (1) and acetone (2). Concentration of perchloric Acid - 2.5.  
KEY: (a) concentration, M.

Increase of  $G(\text{Fe}^{3+})$  during increase of concentration of ions of  $\text{Fe}^{2+}$  occurs due to an increase of fraction of reaction (6) and the corresponding decrease of fraction of reaction (3) and, possibly, as a result of involvement, in the reaction of oxidation of iron, of a certain fraction of excited molecules of water.


On addition of an organic substance to the irradiated solution of inorganic

compound usually is observed an increase of yield of transformation of inorganic compound. Such effects were observed during the irradiation of mixtures of organic compounds with sodium nitrate [6], cerium sulfate [7], sulfate of iron [8], bichromate of potassium [9], et al. During an irradiation of solutions of ethyl alcohol and acetone in 2.5 M  $\text{HClO}_4$  we detected a decrease of yield of reduction of iron of perchlorate that was the greater the higher the concentration of organic compound (Fig. 4). In Fig. 5 is shown the dependence of yield of reduction of ion of perchlorate on the cube root of the concentration of organic compound. In this case the curve of yields of reduction becomes a straight line in the area of concentrations not exceeding 1 M, and is described by the following equations:

$$G(-\text{ClO}_4^-) = 1,25 - (0,61 \pm 0,05) \sqrt[3]{[\text{C}_2\text{H}_5\text{OH}]}$$

$$G(-\text{ClO}_4^-) = 1,25 - (0,32 \pm 0,01) \sqrt[3]{[\text{CH}_3\text{COCH}_3]}$$

In the presence of organic compound the accumulation of no other products of reduction of ion of perchlorate, besides ions of chloride and chlorate is detected; it follows from this that decrease of yields of reduction of ion of perchlorate occurs at the primary stage of decomposition. The observed decrease of yield of reduction probably, is caused by reaction between radicals of perchlorate formed as a result of direct action and molecules of organic compound:



Recently was detected a similar reaction [10,11] between acetonitrile and a radical of perchlorate appearing during anodic oxidation of perchlorate of silver.




Fig. 5. Dependence of yield of disintegration of ion of perchlorate on the cube root of the concentration of ethyl alcohol (1) and acetone (2). Concentration of  $\text{HClO}_4$  - 2.5 M.

One should note that molecules of investigated organic compounds in relation to the radical of perchlorate are more active, than the ion  $\text{Fe}^{2+}$  (for 0.06 M solutions no less than 10 times). This is not surprising, since it is known that in relation to the radical  $\text{OH}$ , which will behave analogously to the radical  $\text{ClO}_4$ , the ion  $\text{Fe}^{2+}$  <sup>is</sup> several times less active than certain organic compounds, for example butyl alcohol [12].

The process of disintegration of radical  $\text{ClO}_4$  by reaction (2) apparently has a very high rate. Therefore it is possible to assume that reaction (2) proceeds in places of high density of ionization, in spurs. Reaction (10), should thus also proceed in spurs. Probability of collision between radical  $\text{ClO}_4$  and the molecule of organic compound will then depend not on the concentration of organic molecules in the entire volume, but on the linear concentration; consequently it will be equal to the cube root of the concentration in the volume. This explains the dependence of  $G(-\text{ClO}_4^-)$  on the cube root of the concentration of acetone and alcohol. The considered dependence is analogous to the dependence of molecular yield of hydrogen on the concentration of sodium nitrate [13] and of molecular hydrogen peroxide on the concentration of chloride and bromide of potassium [14].




Fig. 6. Dependence of ratio of ion yields of chlorate and chlorine ions on the concentration of ethyl alcohol (1) and acetone (2). Concentration of  $\text{HClO}_4$  - 2.5M.

KEY: (a) concentration, M.

In Fig. 6 is shown the change of ratio  $G(\text{ClO}_3^-)/G(\text{Cl}^-)$  during increase of concentration of organic compounds in the solution. On addition of ethyl alcohol and acetone, good acceptors of  $\text{OH}$  radicals, the ratio of yields is decreased. It

is possible to note that the lowering of the ratio of yields occurs significantly more slowly, than in the presence of ions  $\text{Fe}^{2+}$ . This is connected, probably, with the fact that molecules of organic compounds do not reduce the dioxide of chlorine, and the decrease of yield of chlorate occurs due to capture of radicals OH by molecules of organic compounds.

In conclusion I would like to thank V. N. Belevskiy and Wu Tsi-lan for their help in carrying out the experiments, and Professor N. A. Bakh for discussion and valuable advice concerning carrying out of the given work.

M. V. Lomonosov, Moscow State University  
Chemistry Department

#### Literature

1. M. Cottin. J. Chem. Phys., 53, 903 (1956).
2. L. T. Bugayenko and V. N. Belevskiy. Herald of Moscow State University, Chemical Series, No. 2, 77 (1961).
3. L. T. Bugayenko. Herald of Moscow State University, Chemical Series, No. 3, 21 (1961).
4. A. O. Allen. Rad. Res., 1, 85 (1954).
5. T. Lee. Action of radiation in living cells. Cambridge University Press. London, 1946.
6. B. A. Sharpatyy, V. D. Orekhov and M. A. Proskurnin. DAN SSSR, 122, 852 (1958).
7. G. Clark and W. Coe. J. Chem. Phys., 5, 97 (1937).
8. H. Dewhurst. J. Chem. Phys., 19, 1329 (1951).
9. H. Fricke and E. R. Brownscombe. J. Amer. Chem. Soc., 55, 2358 (1933).
10. H. Schmidt and J. Noack. A. anorgan. allg. Chem., 296, 262 (1958).
11. A. H. Maki and D. H. Geske. J. Chem. Phys., 30, 1356 (1959).
12. F. S. Dainton. J. Chem. Phys., 52, 593 (1955).
13. H. A. Mahlman. J. Chem. Phys., 31, 993 (1959).
14. T. S. Sworsky. Rad. Res., 2, 26 (1955).

# DISTRIBUTION LIST

DEPARTMENT OF DEFENSE	Nr. Copies	MAJOR AIR COMMANDS	Nr. Copies
		AFSC	
		TDBTL	5
HEADQUARTERS USAF		TDBDP	10
		TDEPF (Markle)	1
AFGIM-3D2	1	TDGS	1
		TDEMT	10
		TDEY (Honaker)	1
		TDEMT (Endicott)	1
		AEDC (AEY)	1
OTHER AGENCIES		AFWL (WLF)	1
		ASD (ASFA)	2
CIA	5	ESD (ESY)	1
DIA	4		
AID	2		
NASA	1		
AFCLL (CRLLR)	1		
PGE	12		